The Kamil Crater in Egypt

Luigi Folco,[1]* Mario Di Martino,[2] Ahmed El Barkooky,[3] Massimo D'Orazio,[4] Ahmed Lethy,[5] Stefano Urbini,[6] Iacopo Nicolosi,[6] Mahfooz Hafez,[5] Carole Cordier,[1] Matthias van Ginneken,[1] Antonio Zeoli,[1] Ali M. Radwan,[5] Sami El Khrepy,[5] Mohamed El Gabry,[5] Mahomoud Gomaa,[5] Aly A. Barakat,[7] Romano Serra,[8] Mohamed El Sharkawi[3]

Impact craters up to a few hundreds of meters in diameter are common structures of solid surfaces of planetary bodies in the solar system. Statistics predict that impacts producing small craters on Earth occur on decadal to secular time scales (*1*, *2*). However, small craters are rare on Earth because they are rapidly eroded, and the few identified so far [15 <300 m in diameter out of 176 craters up to 300 km in diameter (*3*)] have lost most of their primary features.

We report the detection in southern Egypt of a rayed impact crater 45 m in diameter (Fig. 1A) on a Cretaceous sandstone target. The ejecta rays highlight the exceptional freshness of the structure. The crater was identified by V. De Michele during a Google Earth survey and named Kamil Crater after nearby Gebel Kamil. A geophysical expedition undertaken [supporting online material (SOM)] in February 2010 revealed that the crater is bowl shaped and has an upraised rim (~3 m above preimpact surface) (figs. S1 and S2) typical of simple craters (*4*). The true crater floor depth is 16 m and is overlain by ~6-m-thick crater-fill material (fig. S2). Morphometric parameters agree with those predicted by models (*5*) for a transient crater generated by an iron meteorite 1.3 m in diameter (equivalent to 9.1×10^3 kg) impacting at a velocity of 3.5 km s^{-1}, assuming an average meteoroid entry velocity and entry angle of 18 km s^{-1} and 45°, respectively. Centimeter-scale masses of scoriaceous impact melt glass (fig. S3) occur in and close to the crater and indicate local shock pressures >60 GPa (*4*). We identified 5178 iron meteorite specimens totaling ~1.71 tons in the crater and surrounding area during systematic searches (SOM). They consist of <34-kg shrapnel produced by the explosion of the impactor upon hypervelocity collision with the target (Fig. 1B), except one individual fragment of 83 kg (fig. S4). This evidence indicates that the Kamil Crater was generated by an impactor that landed nearly intact without substantial fragmentation in the atmosphere. The meteorite is classified as an ungrouped Ni-rich ataxite [Ni = 19.8 weight % (wt %), Co = 0.75 wt %, Ga = 49.5 μg g^{-1}, Ge = 121 μg g^{-1}, Ir = 0.39 μg g^{-1}; data following (*6*); fig. S5]. Magnetic anomaly data show no evidence of buried meteorites larger than some tens of centimeters (fig. S1).

On the basis of systematic meteorite searches, the estimated total mass of the impactor is of the order of 5×10^3 to 10×10^3 kg, corresponding to a preatmospheric mass of ~20×10^3 to 40×10^3 kg (*2*). According to geophysical models (*2*, *7*), iron masses <3×10^6 kg normally fragment upon impact with Earth's atmosphere, thereby reducing the energy of the impact at Earth's surface. However, the present statistics, which include the recently discovered Whitecourt Crater (*8*) and the Kamil Crater, suggest that ~35% of the iron meteorites in the above mass range are not disrupted in the atmosphere.

References and Notes

1. P. A. Brown, R. E. Spalding, D. O. ReVelle, E. Tagliaferri, S. P. Worden, *Nature* **420**, 294 (2002).
2. P. A. Bland, N. A. Artemieva, *Meteorit. Planet. Sci.* **41**, 607 (2006).
3. Earth Impact Database, www.unb.ca/passc/ImpactDatabase.
4. H. J. Melosh, *Impact Cratering: A Geologic Process*, Oxford Monographs on Geology and Geophysics (Oxford Univ. Press, Oxford, 1989).
5. G. S. Collins, H. J. Melosh, R. A. Marcus, *Meteorit. Planet. Sci.* **40**, 817 (2005).
6. M. D'Orazio, L. Folco, *Geostand. Newsl.* **27**, 215 (2003).
7. P. A. Bland, N. A. Artemieva, *Nature* **424**, 288 (2003).
8. C. D. K. Herd *et al.*, *Geology* **36**, 955 (2008).
9. Work carried out within the "2009 Egyptian-Italian Year of Science and Technology" framework. T. Hussein and F. Porcelli provided diplomatic support; the Egyptian Army, logistical support; G. Negro, field assistance; F. Speranza and P. Rochette, helpful discussion; the European Commission (Marie Curie Actions–RTNs ORIGINS project), Fondazione Cassa di Risparmio di Torino, Banca Monte dei Paschi di Siena S.p.A., and Telespazio S.p.A. (in the European Space Agency–Space Situational Awareness program framework), financial support.

Supporting Online Material

www.sciencemag.org/cgi/content/full/science.1190990/DC1
Materials and Methods
Figs. S1 to S5

15 April 2010; accepted 29 June 2010
Published online 22 July 2010;
10.1126/science.1190990
Include this information when citing this paper.

[1]Museo Nazionale dell'Antartide Università di Siena, Via Laterina 8, 53100 Siena, Italy. [2]Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Torino, 10025 Pino Torinese, Italy. [3]Department of Geology, Faculty of Sciences, Cairo University, Giza, Egypt. [4]Dipartimento di Scienze della Terra, Università di Pisa, Via S. Maria 53, 56126 Pisa, Italy. [5]National Research Institute of Astronomy and Geophysics, Helwan, Egypt. [6]Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata 605, 00143 Roma, Italy. [7]Egyptian Mineral Resources Authority, 3 Salah Salem Road, Abassiya, Cairo, Egypt. [8]Dipartimento di Fisica, Università di Bologna, Via Irnerio 46, 40126 Bologna, Italy.

*To whom correspondence should be addressed. E-mail: folco@unisi.it




Fig. 1. (**A**) QuickBird satellite image (22 October 2005; courtesy of e-GEOS) of Kamil Crater. (**B**) An ~3-kg shrapnel of the associated iron meteorite.

Downloaded from www.sciencemag.org on August 13, 2010